



OCA2 gene

OCA2 melanosomal transmembrane protein

Normal Function

The *OCA2* gene (formerly called the *P* gene) provides instructions for making a protein called the P protein. This protein is located in melanocytes, which are specialized cells that produce a pigment called melanin. Melanin is the substance that gives skin, hair, and eyes their color. Melanin is also found in the light-sensitive tissue at the back of the eye (the retina), where it plays a role in normal vision.

Although the exact function of the P protein is unknown, it is essential for normal pigmentation and is likely involved in the production of melanin. Within melanocytes, the P protein may transport molecules into and out of structures called melanosomes (where melanin is produced). Researchers believe that this protein may also help regulate the relative acidity (pH) of melanosomes. Tight control of pH is necessary for most biological processes.

Health Conditions Related to Genetic Changes

Angelman syndrome

The *OCA2* gene is located in a region of chromosome 15 that is often deleted in individuals with Angelman syndrome. A loss of this gene does not cause the characteristic neurologic features of Angelman syndrome; however, people with this condition who are missing one copy of the *OCA2* gene tend to have unusually light-colored hair and fair skin. Cells with only one copy of the *OCA2* gene make a reduced amount of P protein compared with cells with two functional copies of this gene, which affects the coloring of the hair and skin.

A small percentage of people with Angelman syndrome also have oculocutaneous albinism type 2. This condition occurs when people have two nonfunctional copies of the *OCA2* gene in each cell. In addition to a deletion in chromosome 15 that removes one copy of the *OCA2* gene, these individuals have a mutation in the *OCA2* gene on the other copy of chromosome 15. As a result, cells make little or no functional P protein. A lack of P protein disrupts the production of melanin, leading to the characteristic features of albinism.

oculocutaneous albinism

More than 80 mutations in the *OCA2* gene have been identified in people with oculocutaneous albinism type 2. People with this form of albinism often have light yellow, blond, or light brown hair; creamy white skin; light-colored eyes; and problems

with vision. The most common *OCA2* mutation is a large deletion in the gene, which is found in many affected individuals of sub-Saharan African heritage. Other *OCA2* gene mutations, including changes in single DNA building blocks (base pairs) and small deletions, are more common in other populations. Mutations in the *OCA2* gene disrupt the normal production of melanin, which reduces coloring of the hair, skin, and eyes and affects vision.

Prader-Willi syndrome


The region of chromosome 15 containing the *OCA2* gene is often deleted in individuals with Prader-Willi syndrome. A loss of this gene does not cause intellectual disability and the other characteristic features of Prader-Willi syndrome; however, people with this condition who are missing one copy of the *OCA2* gene tend to have unusually light-colored hair and fair skin. Cells missing a copy of the *OCA2* gene make less P protein than cells with two functional copies of the gene, which affects the coloring of the hair and skin.

Oculocutaneous albinism type 2 also occurs in a small number of people with Prader-Willi syndrome. This condition occurs when people have two nonfunctional copies of the *OCA2* gene in each cell. In addition to a deletion in chromosome 15 that removes one copy of the *OCA2* gene, these individuals have a mutation in the *OCA2* gene on the other copy of chromosome 15. As a result, cells make little or no functional P protein. A lack of P protein disrupts the production of melanin, leading to the characteristic features of albinism.

Chromosomal Location

Cytogenetic Location: 15q12-q13.1, which is the long (q) arm of chromosome 15 between positions 12 and 13.1

Molecular Location: base pairs 27,719,008 to 28,099,342 on chromosome 15 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- BOCA
- Melanocyte-specific transporter protein
- oculocutaneous albinism II
- oculocutaneous albinism II (pink-eye dilution homolog, mouse)
- P gene
- P_HUMAN
- PED
- Pink-eyed dilution protein homolog

Additional Information & Resources

Educational Resources

- Protein Spotlight, Swiss Institute of Bioinformatics
http://web.expasy.org/spotlight/back_issues/054/

GeneReviews

- Angelman Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1144>
- Oculocutaneous Albinism Type 2
<https://www.ncbi.nlm.nih.gov/books/NBK1232>
- Prader-Willi Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1330>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28OCA2%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- OCA2 GENE
<http://omim.org/entry/611409>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
<http://atlasgeneticsoncology.org/Genes/OCA2ID45789ch15q12.html>
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=OCA2%5Bgene%5D>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=8101
- International Albinism Center, University of Minnesota: Albinism Database
<http://www.ifpcs.org/albinism/oca2mut.html>
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/4948>
- UniProt
<http://www.uniprot.org/uniprot/Q04671>

Sources for This Summary

- Bittel DC, Kibiryeva N, Talebizadeh Z, Butler MG. Microarray analysis of gene/transcript expression in Prader-Willi syndrome: deletion versus UPD. J Med Genet. 2003 Aug;40(8):568-74.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12920063>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1735542/>
- Bittel DC, Kibiryeva N, Talebizadeh Z, Driscoll DJ, Butler MG. Microarray analysis of gene/transcript expression in Angelman syndrome: deletion versus UPD. Genomics. 2005 Jan;85(1):85-91.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15607424>
- Brilliant MH. The mouse p (pink-eyed dilution) and human P genes, oculocutaneous albinism type 2 (OCA2), and melanosomal pH. Pigment Cell Res. 2001 Apr;14(2):86-93. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11310796>
- Duffy DL, Montgomery GW, Chen W, Zhao ZZ, Le L, James MR, Hayward NK, Martin NG, Sturm RA. A three-single-nucleotide polymorphism haplotype in intron 1 of OCA2 explains most human eye-color variation. Am J Hum Genet. 2007 Feb;80(2):241-52. Epub 2006 Dec 20.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17236130>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1785344/>
- Fridman C, Hosomi N, Varela MC, Souza AH, Fukai K, Koiffmann CP. Angelman syndrome associated with oculocutaneous albinism due to an intragenic deletion of the P gene. Am J Med Genet A. 2003 Jun 1;119A(2):180-3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12749060>
- GeneReview: Oculocutaneous Albinism Type 2
<https://www.ncbi.nlm.nih.gov/books/NBK1232>
- Kerr R, Stevens G, Manga P, Salm S, John P, Haw T, Ramsay M. Identification of P gene mutations in individuals with oculocutaneous albinism in sub-Saharan Africa. Hum Mutat. 2000;15(2):166-72. Erratum in: Hum Mutat 2000;16(1):following 86.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10649493>

- Oetting WS, Garrett SS, Brott M, King RA. P gene mutations associated with oculocutaneous albinism type II (OCA2). Hum Mutat. 2005 Mar;25(3):323.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15712365>
- Saitoh S, Oiso N, Wada T, Narazaki O, Fukai K. Oculocutaneous albinism type 2 with a P gene missense mutation in a patient with Angelman syndrome. J Med Genet. 2000 May;37(5):392-4.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10905897>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1734585/>
- Spritz RA, Bailin T, Nicholls RD, Lee ST, Park SK, Mascari MJ, Butler MG. Hypopigmentation in the Prader-Willi syndrome correlates with P gene deletion but not with haplotype of the hemizygous P allele. Am J Med Genet. 1997 Jul 11;71(1):57-62.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/9215770>
- Suzuki T, Miyamura Y, Matsunaga J, Shimizu H, Kawachi Y, Ohyama N, Ishikawa O, Ishikawa T, Terao H, Tomita Y. Six novel P gene mutations and oculocutaneous albinism type 2 frequency in Japanese albino patients. J Invest Dermatol. 2003 May;120(5):781-3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12713581>
- Toyofuku K, Valencia JC, Kushimoto T, Costin GE, Virador VM, Vieira WD, Ferrans VJ, Hearing VJ. The etiology of oculocutaneous albinism (OCA) type II: the pink protein modulates the processing and transport of tyrosinase. Pigment Cell Res. 2002 Jun;15(3):217-24. Erratum in: Pigment Cell Res. 2002 Oct;15(5):400..
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12028586>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/OCA2>

Reviewed: October 2011
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services